



EASTERN UNIVERSITY, SRI LANKA
DEPARTMENT OF MATHEMATICS
FIRST EXAMINATION IN SCIENCE, 2010/2011
FIRST SEMESTER (Nov./Dec., 2012)
AM 106 - TENSOR CALCULUS
(Proper & Repeat)

Answer all questions

Time : One hour

1. (a) Explain what is meant by the following terms:
 - i. Covariant tensor;
 - ii. Contravariant tensor.
- (b) Write down the law of transformation for the following tensors:
 - i. A_{mn} ;
 - ii. B_r^{pq} ;
 - iii. C_{rt}^{pqs} .
- (c) If $ds^2 = g_{ij}dx^i dx^j$ is an invariant, show that g_{ij} is a symmetric covariant tensor of rank two.
- (d) Express the relationship between the following associated tensors:
 - i. A^{jkl} and A_{pqr} ;
 - ii. A_j^k and A^{kr} .

(e) If $X(i, j) B^j = C_i$, where B^j is an arbitrary contravariant vector and C_i is a covariant vector, then show that $X(i, j)$ is a tensor. What is its rank and type.

2. (a) Define the following:

- i. Christoffel's symbols of the first and second kind;
- ii. Geodesic;
- iii. Covariant derivative of A_p .

(b) With the usual notations, prove the following:

- i. $[pq, r] = g_{rs} \Gamma_{pq}^s$;
- ii. $[pm, q] + [qm, p] = \frac{\partial g_{pq}}{\partial x^m}$;
- iii. $\frac{\partial g^{pq}}{\partial x^m} + g^{pn} \Gamma_{mn}^q + g^{qn} \Gamma_{mn}^p = 0$.

Hence show that,

$$g_{jk;q} = 0.$$

(c) Show that the non-vanishing Christoffel's symbols of the second kind in cylindrical coordinate (ρ, ϕ, z) are given by

$$\Gamma_{22}^1 = -\rho, \quad \Gamma_{21}^2 = \frac{1}{\rho}, \quad \Gamma_{12}^2 = \frac{1}{\rho},$$

where $x^1 = \rho$, $x^2 = \phi$, $x^3 = z$.