



EASTERN UNIVERSITY, SRI LANKA  
DEPARTMENT OF MATHEMATICS

EXTERNAL DEGREE EXAMINATION IN SCIENCE - 2009/2010

SECOND YEAR, FIRST SEMESTER (June/Sept., 2012)

EXTMT 207 - NUMERICAL ANALYSIS

(PROPER & REPEAT)

---

Answer all questions

Time: Two hours

---

- Q1. (a) Write the suitable form of any non-zero number  $x \in F$ , where  $F$  represents the set of all floating point numbers, and identify the terms involved.
- (b) Define the relative round-off error, and explain with an illustrative example.
- (c) Find the absolute and relative errors if the computed answer of the exact value 10.147 is 10.159.
- (d) A function  $f(x) = x^3 - 3x^2 + 3x - 1$  is rearranged in a nested form given by

$$g(x) = [(x - 3)x + 3]x - 1.$$

Find  $f(2.19)$  and  $g(2.19)$  using 3-digit rounding. If the true value of  $f(x)$  and  $g(x)$  at  $x = 2.19$  is 1.685159, compare the errors, and state the significance of this problem.

- Q2. (a) (i) Let  $x = g(x)$  be an arrangement of an equation  $f(x) = 0$ , which has a root  $\alpha$  in the interval  $I$ . Suppose that  $g'(x)$  exists and is continuous in  $I$  such that

$$|g'(x)| \leq h < 1, \forall x \in I,$$

where  $0 < h < 1$ .

Prove that for any given  $x_0$ , the sequence  $\{x_r\}$ ,  $r = 0, 1, 2, \dots$ , defined by

$$x_{r+1} = g(x_r)$$

converges to the root  $\alpha$ , and such  $\alpha$  is unique.

- (ii) Following iterative formulas are proposed to find a real root of the equation  $f(x) = x^3 + x^2 - 1 = 0$ , using the iterative method given in (i).

$$x_{r+1} = \frac{1}{\sqrt{x_r + 1}}$$
$$x_{r+1} = \frac{1}{x_r^2} - 1.$$

Check the applicability of iterative equations (1) and (2) in finding real root of  $f(x)$ .

- (b) Derive the Newton-Raphson method using Taylor series or otherwise. Carry out four iterations to find  $x$ , correct to 4-decimal points, such that

$$f(x) = x^4 - 5 = 0$$

with an initial estimate  $x_0 = 2$ .

- Q3. (a) Write down the divided difference table for  $e^x$  using the values

$x$	$e^x$
0.0	1.00000
0.4	1.49182
0.9	2.45960
1.5	4.48169
1.8	6.04965

Estimate  $e^{1.2}$ , correct to 4-decimal places, using second and third degree interpolation polynomials. If the exact value of  $e^{1.2}$  is 3.3201, which interpolation polynomial gives the better estimate? Justify your answer.

- (b) Use the Composite Trapezium rule with 2, 4 and 8 sub-intervals to estimate the integral

$$I = \int_1^2 e^x dx.$$

If the exact value of  $I$  is 4.67078, tabulate the error in each case. What do you say about the accuracy with respect to step size?



Q4. (a) Solve the system of equations

$$\begin{aligned}4x_1 + 4x_2 + x_3 + 4x_4 &= 12 \\2x_1 + 5x_2 + 7x_3 + 4x_4 &= 1 \\10x_1 + 5x_2 - 5x_3 &= 25 \\-2x_1 - 2x_2 + x_3 - 3x_4 &= -10\end{aligned}$$

using the Gaussian elimination.

(b) Solve the system of equations

$$\begin{aligned}16x_1 - 4x_2 + 4x_3 &= 24 \\-4x_1 + 5x_2 + 3x_3 &= -6 \\4x_1 + 3x_2 + 14x_3 &= 15\end{aligned}$$

by applying the Jacobi iteration (complete 3 iterations with rounding correct to 4-decimal points) using the initial guess  $x_1^{(0)} = 0$ ,  $x_2^{(0)} = 0$  and  $x_3^{(0)} = 0$ .