

**NUTRITIONAL ASSESSMENT OF
PRE - SCHOOL CHILDREN IN PADDIPALAI
DIVISIONAL SECRETARIAT AREA IN
BATTICALOA DISTRICT**



By

SUBRAMANIYAM SISIKARAN



FAG330



Project Report
Library - EUSL

FACULTY OF AGRICULTURE

EASTERN UNIVERSITY

SRILANKA

2011

**PROCESSED
Main Library, EUSL**

ABSTRACT

In Sri Lanka, state of malnutrition among children is unfavorable. The situation may worse in Batticaloa District where civil war exists. Therefore, this study was conducted to assess nutritional assessment of pre-school children in Paddipalai Divisional Secretariat area in Batticaloa District. A sample of 150 subjects was selected from pre-schools through multi stage sampling techniques. Anthropometric measurements were performed using standardized procedure on the preschool children, Weight, Height, Mid Upper Arm Circumference, Hip and Waist were measured. Anthropometric indices such as Height-for-age, Weight-for-height, and Weight-for-age were generated. Information on socio-economic characteristic of household and dietary pattern of children was gathered by an interview schedule, which includes food frequency questionnaire.

Children showed a considerable risk for being stunted (65%), wasting (61%) and under weight (80%). Girls had higher risk than that of boys. Prevalence of malnutrition is higher in all inter displaced people.

Among the tested individuals all families had been displaced due to severe civil war. Among the tested subjects, 70% of children's families receive food aids or food stamps from government. Most of their parent's income level was under the poverty level of Rs.3000 per month. About 72.2% of the parent's education level is below the G.C.E (O/L) and rate of literacy of them around 27%. The student's frequency of intake of meat is monthly basis and they consume egg, milk and fruit at fortnightly basis. Nutritional status of pre-school children in the Paddipalai Divisional Secretariat in Batticaloa district is very poor and civil unrest, unfavorable climatic changes also to be associated with malnutrition of children.

TABLE OF CONTENT

| | |
|--|-----------|
| ABSTRACT..... | I |
| ACKNOWLEDGEMENTS | II |
| TABLE OF CONTENT | III |
| LIST OF TABLES | VIII |
| LIST OF FIGURES | IX |
| LIST OF ABBREVIATIONS..... | X |
| CHAPTER I..... | 01 |
| 1.0 INTRODUCTION | 01 |
| 1.1 Objectives..... | 06 |
| CHAPTER II..... | 07 |
| 2.0 LITERATURE REVIEW | 07 |
| 2.1 Background information of study area..... | 07 |
| 2.1.1 Geographical map and profile of study area | 07 |
| 2.1.1.1 Major soils in the Division..... | 10 |
| 2.1.2 Population and Economy | 10 |
| 2.1.2.1 Population | 10 |
| 2.1.1.1 Economy | 11 |
| 2.1.3 Status of Industries..... | 13 |
| 2.1.4 Housing and Infrastructure | 13 |
| 2.1.5 Status of Fishery..... | 13 |
| 2.1.6 Status of Agriculture | 14 |
| 2.1.7 Education | 15 |
| 2.1.8 Infrastructure Facilities..... | 16 |
| 2.1.8.1 Electricity..... | 16 |
| 2.1.8.2 Postal service..... | 16 |

| | |
|---|----|
| 2.1.8.3 Roads and Transport | 16 |
| 2.1.9 Status of Health and Nutrition | 17 |
| 2.2 Survey studies | 18 |
| 2.2.1 Survey on Pre-school children..... | 18 |
| 2.2.2 Diet consumption studies..... | 19 |
| 2.3 Methods of Evaluating state of Malnutrition | 19 |
| 2.3.1 Anthropometric Data..... | 19 |
| 2.3.1.1 Weight..... | 20 |
| 2.3.1.2 Height..... | 21 |
| 2.3.1.3 Weight for Height | 22 |
| 2.3.1.4 Mid Upper Arm Circumference (MUAC) | 22 |
| 2.3.1.5 Skin Fold Thickness..... | 23 |
| 2.3.2 Dietary Assessment..... | 23 |
| Inquiry or Recall..... | 24 |
| Combined observation and inquiry | 25 |
| 2.3.2.1 Food Intake record | 26 |
| 2.3.2.2 Diet History..... | 26 |
| 2.3.2.3 Recall of 24-hour | 27 |
| 2.3.2.4 Weighted Food Intake | 27 |
| 2.3.3 Clinical Examinations | 27 |
| 2.3.4 Biochemical Analysis | 28 |
| 2.4 Protein Energy Malnutrition (PEM) | 28 |
| Under Nutrition | 29 |
| Over Nutrition | 30 |
| School Children..... | 30 |
| 2.4.1 Kwashiorkor..... | 32 |

| | |
|--|-----------|
| 2.4.1.1 The nature of Kwashiorkor disease..... | 33 |
| 2.4.1.1.1 Mental changes | 33 |
| 2.4.1.1.2 Hair changes..... | 33 |
| 2.4.1.1.3 Skin changes | 34 |
| 2.4.1.1.4 Diarrhoea..... | 34 |
| 2.4.1.1.5 Anemia..... | 34 |
| 2.4.2 Marasmus..... | 35 |
| 2.4.2.1 The nature of Marasmus Disease | 36 |
| 2.5 Natural History of Stunting and Wasting..... | 37 |
| 2.5.1 Major nutritional problem in Sri Lanka | 38 |
| 2.5.1.1 Protein Energy Malnutrition | 38 |
| 2.5.1.2 Micro nutrient Malnutrition in Sri Lanka | 39 |
| 2.5.1.2.1 Vitamin A deficiency..... | 39 |
| 2.5.1.2.2 Iron (Fe) deficiency Anemia | 40 |
| 2.5.1.2.3 Iodine (I_2) deficiency disorders..... | 40 |
| CHAPTER III | 41 |
| 3.0 METHODOLOGY..... | 41 |
| 3.1 Sampling Methods and Techniques | 41 |
| 3.2 Data collection and Techniques | 41 |
| 3.2.1 Pilot study | 42 |
| 3.2.2 Definitive study | 42 |
| 3.3 Collecting Data and Anthropometric Measurements..... | 42 |
| 3.3.1 Anthropometric Meaurements | 42 |
| 3.3.1.1 Weight..... | 43 |
| 3.3.1.2 Height..... | 44 |
| 3.3.1.3 Mid Upper Arm Circumference (MUAC) | 44 |

| | |
|---|-----------|
| 3.3.1.4 Skin fold thickness | 45 |
| 3.3.1.5 Waist and Hip circumferences | 46 |
| 3.3.2 Health Status | 47 |
| 3.3.3 Collecting Dietary Information..... | 47 |
| 3.3.4 Gathering the socio-economic details of Parents or Guardians | 48 |
| 3.4 Calculating Indicators and used cut off value | 48 |
| 3.4.1 Body Mass Index (BMI) | 48 |
| 3.4.2 Stunting | 49 |
| 3.4.3 Wasting | 49 |
| 3.4.4 Mid Upper Arm Circumference (MUAC) and Tricep Skin Fold Thickness (TSF)..... | 50 |
| 3.4.5 Waist (cm)/Hip (cm) ratio..... | 50 |
| 3.4.6 Waist (cm)/Height (cm) ratio | 50 |
| 3.5 Pre-Testing | 51 |
| 3.6 Analysis..... | 51 |
| 3.7 Calculation of 'Z' scores..... | 51 |
| CHAPTER IV..... | 52 |
| 4.0 RESULTS AND DISCUSSION | 52 |
| 4.1 Nutritional status of Children..... | 52 |
| 4.1.1 Anthropometric Characteristics | 52 |
| 4.1.1.1 Body Mass Index (BMI) | 52 |
| 4.1.1.2 Distribution of State of Malnutrition | 53 |
| 4.1.1.3 Prevalence of Stunting | 54 |
| 4.1.1.4 Wasting | 55 |
| 4.1.1.5 Prevalence of Under-weight..... | 56 |
| 4.1.1.6 Mid Upper Arm Circumference..... | 57 |

| | |
|---|-----------|
| 4.1.1.7 Waist Hip ratio..... | 58 |
| 4.1.1.8 Waist Height ratio | 58 |
| 4.2 Health status..... | 59 |
| 4.3 Frequency of Food intake..... | 60 |
| 4.4 General Living situations..... | 61 |
| 4.4.1 Household characters | 61 |
| 4.4.2 Displacement..... | 61 |
| 4.4.3 Education of Father and Mother | 62 |
| 4.4.4 Income sources..... | 63 |
| 4.5 Infectious Diseases..... | 64 |
| 4.6 Dependency of Family..... | 65 |
| 4.7 Sanitary Practices..... | 66 |
| 4.8 Relationship between factors Affecting Nutrition Status. | 67 |
| 4.8.1 Monthly Family Income and BMI value | 67 |
| 4.8.2 Number of Family members and BMI value | 68 |
| 4.8.3 Education level of Parents and BMI value..... | 68 |
| 4.9 General Comments..... | 69 |
| 4.10 Limitation of the study..... | 70 |
| CHAPTER V | 71 |
| 5.0 CONCLUSION | 71 |
| SUGGESTIONS | 73 |
| REFERENCES..... | 75 |
| APPENDICES | 81 |