DEVELOPMENT AND QUALITY EVALUATION OF NUTRITIONALLY ENRICHED COMPLEMENTARY FOOD FOR INFANTS



5

BY

ROSHANA MOHAMED RASHEED





DEPARTMENT OF AGRICULTURAL CHEMISTRY FACULTY OF AGRICULTURE

EASTERN UNIVERSITY

SRI LANKA

2017

PROCESSED Main Library, EUSL

DEVELOPMENT AND QUALITY EVALUATION OF NUTRITIONALLY ENRICHED COMPLEMENTARY FOOD FOR INFANTS



2

BY

ROSHANA MOHAMED RASHEED





DEPARTMENT OF AGRICULTURAL CHEMISTRY

FACULTY OF AGRICULTURE

EASTERN UNIVERSITY

SRI LANKA

2017

PROCESSED Main Library, EUSI

ABSTRACT

Protein-energy malnutrition and micronutrient deficiency among children are the major health challenges in developing countries. It affects the child at the most crucial period, *i.e.* stage of development, which can lead to permanent impairment in later life. Usually cereals based fortified foods are introduced initially as the first complementary foods at the six months of infancy. Although, a number of convenient cereal formulas are available, they are often too expensive for the poor and middle income people. To overcome this problem present study was designed to aim at the formulation of low cost complementary foods that provide sufficient protein, vitamin and minerals for children.

The complementary food mixtures were prepared from unpolished parboiled red rice flour, germinated green gram flour and carrot flour in the ratios of 100:00:00, 80:10:10, 70:20:10, 60:30:10, 50:40:10 and 40:50:10 respectively. The mixtures were subjected to nutritional (moisture, ash, protein, fat, fiber and carotene), organoleptic (colour, texture, taste, aroma and overall acceptability) and microbial analysis to evaluate the suitability of these complementary food mixtures for consumption and for long shelf life. The nutritional analysis of the complementary food mixture revealed that the moisture, ash, protein and fiber content increased and fat and vitamin A decreased with the increasing of the germinated green gram flour from 10 to 50%. The sensory assessment showed that, there were significant differences (p<0.05) among the treatments.

Based on the quality characteristics, most preferred complementary food mixtures were selected and stored for 14 weeks at 30±1°C and 75-80% RH. The results of storage studies showed the declining trends in ash, protein, fiber, fat and vitamin A and an

i

increasing trend in moisture of the complementary food mixture. The results of nutritional analysis showed that, there were significant differences (p<0.05) among the treatments. From the overall acceptability rating, the complementary food mixture with 60% unpolished parboiled red rice flour, 30% germinated green gram flour and 10% carrot flour had the highest mean value compared with other tested treatments. There were no total plate counts observed in the formulated complementary food mixture during the entire storage period. The results indicated that the complementary food mixture flour and 10% carrot flour contained 14.2% protein, 1.6% fiber, 2.06% ash, 1.91% fat, 6.28% moisture and 30.23 mg kg⁻¹ vitamin A, and which could be stored for 14 weeks without any significant changes in the quality characteristics.

ii

1

TABLE OF CONTENTS

ABSTRACTi
ACKNOWLEDGEMENTiii
TABLE OF CONTENTSiv
LIST OF TABLESxii
LIST OF FIGURESxiii
LIST OF PLATESxiv
CHAPTER 01 - INTRODUCTION1
CHAPTER 02 - LITERATURE REVIEW
2.1 Nutrition
2.1.1 Nutrients
2.1.1.1 Carbohydrates
2.1.1.2 Proteins
2.1.1.3 Fats
2.1.1.4 Vitamins
2.1.1.5 Minerals
2.1.1.6 Water
2.2 Major Nutritional Problems in Sri Lanka
2.2.1 Malnutrition
2.2.1.1 Protein-Energy Malnutrition (PEM)10
2.2.1.2 Micronutrient Deficiencies10
2.2.1.2.1 Vitamin A Deficiency11
2.2.2 Nutritional Status in Sri Lanka

2.3 Complementary Feeding
2.3.1 Definition
2.3.2 Complementary Time
2.3.3 Recommendation of Complementary Food for Infants
2.3.4 Nutritional Requirements of Infants and Problems during Weaning14
2.3.5 Ingredients Used in Complementary Foods15
2.4 Rice
2.4.1 Scientific Classification
2.4.2 Description of Rice
2.4.3 Parboiling of Rice
2.4.4 Nutritional Facts of Rice
2.4.5 Medicinal Uses of Rice
2.5 Green Gram
2.5.1 Scientific Classification
2.5.2 Origin, Domestication and Adaptation
2.5.3 Morphological Description
2.5.4 Nutritional Value of Green Gram
2.5.5 Utilization of Green Gram
2.5.6 Germinated Green Gram
2.5.6.1 Germination Process
2.5.6.2 Nutritional Improvement during Germination23
2.5.6.3 Health Benefits of Germinated Green gram24
2.6 Carrot

2.6.1 Scientific Classification	4
2.6.2 Description of Carrot	5
2.6.3 Nutritional Value of Carrot	5
2.6.4 Morphology of the Carrot2	7
2.6.5 Processing and Use of Carrot	7
2.7 Technology of Roasting	8
2.8 Packaging and Shelf life of Products	9
2.8.1 Packaging Material	9
2.8.2 Laminated Packaging Films	0
2.9 Quality Parameters of Dried and Related Food	1
2.10 Microbiological Examination	1
2.10.1 Total Plate Count	2
2.11 Sensory Evaluations	2
2.11.1 Definition	2
2.11.2 Uses of Sensory Evaluation	3
2.11.3 Problem Associated with Sensory Evaluation	3
2.11.4 Rules of Sensory Evaluation	4
2.11.5 Sensory Evaluation Methods	
2.11.5.1 Discrimination Analysis	5
2.11.5.2 Descriptive Analysis	5
2.11.5.3 Affective Analysis	5
2.11.5.3.1 Hedonic Rating Test	5
2.11.6 Common Practices when Performing Sensory Evaluation Tests	7

1

2.11.6.1 Test Controls
2.11.6.2 Product Controls
2.11.6.3 Panel Controls
2.11.7 Qualities Assessed by Sensory Evaluation
2.11.7.1 Colour
2.11.7.2 Taste
2.11.7.3 Aroma
2.11.7.4 Texture
2.11.7.5 Overall Acceptability
CHAPTER 03 - MATERIALS AND METHODS41
3.1 Materials
3.1.1 Materials Used for the Study41
3.1.2 Material Collection
3.2 Methods
3.2.1 Preparation of Raw Materials for the Formulation of Complementary Food 42
3.2.1.1 Preparation of Unpolished Parboiled Red Rice Flour
3.2.1.2 Preparation of Germinated Green Gram Flour
3.2.1.3 Preparation of Carrot Flour44
3.2.2 Preparation of Instant Complementary Food45
3.2.3 Tested Treatments
3.3 Sensory Evaluation46
3.3.1 Materials Used for the Sensory Evaluation47
3.3.2 Coding the Samples

1

3.3.3 Presentation of the Samples47
3.3.4 Instructions for the Panelists
3.4 Nutritional Analysis of Formulated Complementary Food
3.4.1 Determination of Moisture Content
3.4.1 Determination of 49
3.4.1.2 Materials
3.4.1.2 Materials
3.4.1.4 Calculation
3.4.2 Determination of Ash Content
3.4.2.1 Principle
3.4.2.2 Materials
3.4.2.3 Procedure
3.4.2.4 Calculation
3.4.3 Determination of Protein Content
3.4.3.1 Principle
3.4.3.2 Materials
3.4.3.3 Procedure
3.4.3.4 Calculation
3.4.4 Determination of Fat Content53
3.4.4.1 Principle
3.4.4.2 Materials
3.4.4.3 Procedure
3.4.4.3 Procedure
1.4.4.4 Calculation

62
.3 Quality Characteristics of Freshly Made Complementary Food Mixtures62
4.3.1 Nutritional Qualities of Freshly Made Complementary Food Mixtures62
4.3.1.1 Moisture Content
4.3.1.2 Ash Content
4.3.1.3 Protein Content
4.3.1.4 Fat Content
4.3.1.5 Fiber Content
4.3.1.6 Carotene Content
4.3.2 Organoleptic Evaluation of Complementary Food Mixture
4.3.2.1 Colour
4.3.2.2 Texture
4.3.2.3 Taste
4.3.2.4 Aroma
, 4.3.2.5 Overall Acceptability
4.4 Changes in Quality Characteristics of Complementary Food Mixtures
during Storage at Ambient Temperature
4.4.1 Nutritional Qualities of Complementary Food Mixtures during
Storage
4.4.1.1 Moisture Content71
4.4.1.2 Ash Content72
4.4.1.3 Protein Content73
4.4.1.4 Fat Content74
4.4.1.5 Fiber Content75