

## EASTERN UNIVERSITY, SRI LANKA DEPARTMENT OF MATHEMATICS FIRST EXAMINATION IN SCIENCE -2009/2010 SECOND SEMESTER (April/May, 2012) MT 102 - REAL ANALYSIS (REPEAT)

Answer all Questions

Time: Three hours

10 JUN 2013

- (a) Define the terms Supremum and Infimum of a bounded subset A of ℝ.
   [10marks]
  - (b) Prove that an upper bound u of a non-empty set S in ℝ is the supremum of S if, and only if, for each ε > 0 there exists x<sub>0</sub> ∈ S such that u − ε < x<sub>0</sub>.

[30marks]

- (c) State the Archimedian principle and use it to prove that there exists a positive real number x such that  $x^2 = 2$ . [40marks]
- (d) Use the Mathematical induction principle to show that  $1^2 + 2^2 + \dots + n^2 = \frac{1}{6}n(n+1)(2n+1)$  for all  $n \in \mathbb{N}$ . [20marks]
- (a) Define what is meant by the following terms applied to a sequence of real numbers:
  - i. bounded;
  - ii. convergent;
  - iii. monotone.

[15marks]

- (b) Prove that every increasing sequence of real numbers which is bounded about is convergent. [35mark]
- (c) Let  $(y_n)$  be a sequence of real numbers defined inductively by

$$y_1 = 1,$$
  $y_{n+1} = \frac{1}{4}(2y_n + 3)$  for all  $n \in \mathbb{N}$ .  
Show that  $(y_n)$  is convergent and  $\lim_{n \to \infty} y_n = \frac{3}{2}$ . [50mark]

3. (a) i. Let  $f : \mathbb{R} \to \mathbb{R}$  be a function. Explain what is meant by the function has a limit  $l \in \mathbb{R}$  at a point  $a \in \mathbb{R}$ . [15mark]

ii. Use the definition of the limit to show that  $\lim_{x \to -1} \frac{x+5}{2x+3} = 4$ .

25mark

- (b) i. Let A ⊆ ℝ and f : A → ℝ be a function. Let a ∈ ℝ. Prove th lim f(x) = l exists finitely if, and only if, for every sequence (x<sub>n</sub>) in that converges to a such that x<sub>n</sub> ≠ a for all n ∈ N, the sequence (f(x<sub>i</sub> converges to l. [40mark]
  - ii. Let  $f : \mathbb{R} \setminus \{0\} \to \mathbb{R}$  be defined by  $f(x) = \sin(1/x) \quad \forall x \neq 0$ . Show th  $\lim_{x \to 0} f(x)$  does not exists in  $\mathbb{R}$ . [20mark]
- 4. (a) i. Define what is meant by the statement that a function  $f : \mathbb{R} \to \mathbb{R}$ continuous at a point  $a \ (\in \mathbb{R})$ . [15mark]
  - ii. Show that the function  $f : \mathbb{R} \to \mathbb{R}$  defined by  $f(x) = \cos x$ ,  $\forall x \in \mathbb{R}$ continuous at every point in  $\mathbb{R}$ . [25mark]
  - (b) Let I = [a, b] be a closed and bounded interval in  $\mathbb{R}$ . Prove that if  $f: I \rightarrow$  is continuous on I then f is bounded on I.
  - (c) State the Intermediate Value Theorem and use it to prove that the equation  $2x^2(x+2) 1 = 0$  has a root in each of the intervals (-2, -1), (-1, 0) at (0,1).
- 5. (a) i. Define what is meant by a function  $f : \mathbb{R} \to \mathbb{R}$  is differentiable at the part  $x_0 \in \mathbb{R}$ . [15mark]

ii. Discuss differentiability of each of the following functions  $f : \mathbb{R} \to \mathbb{R}$  at the origin:

1. 
$$f(x) = \sin x$$
  
2.  $f(x) = |x|$   
3.  $f(x) = \begin{cases} 3+x, & x \le 0; \\ 3-x, & x > 0. \end{cases}$  [30marks]

(b) i. Let f: [a, b] → ℝ be a function where a, b ∈ ℝ with a < b. Suppose that f is continuous on [a, b] and differentiable on (a, b). Prove that there exists c ∈ (a, b) such that</li>

$$f'(c) = \frac{f(b) - f(a)}{b - a}$$

(You may use the Rolle's Theorem without proving it.) [30marks] ii. Show that  $x < \sin^{-1} x < \frac{x}{\sqrt{1-x^2}}$   $\forall x \in (0,1)$ . [25marks]

(a) Suppose that f and g are two continuous real valued functions defined on [a, b], where a, b ∈ ℝ with a < b. Suppose also that f and g are differentiable on (a, b) and g'(x) ≠ 0 ∀x ∈ (a, b). Prove that for some c ∈ (a, b),</li>

$$\frac{f'(c)}{g'(c)} = \frac{f(b) - f(a)}{g(b) - g(a)}.$$

(You may use the Rolle's Theorem without proving it.) [30marks]

- (b) i. Suppose that f and g are continuous on [a, b], differentiable on (a, b) and let f(c) = g(c) = 0 for some c ∈ (a, b). Further suppose that g(x) ≠ 0 and g'(x) ≠ 0 for all x ∈ (a, b) \ {c}. If lim f'(x) = l exists finitely prove that lim f(x) = l. [20marks]
  ii. Prove that lim (1 cos x) / x<sup>2</sup> = 1/2. [15marks]
- (c) State the Taylor's Theorem and use it to prove that

$$1 - \frac{1}{2}x^2 \le \cos x \quad \forall x \in \mathbb{R}.$$

[35marks]