EASTERN UNIVERSITY, SRI LANKA

THIRD EXAMINATION IN SCIENCE - 2002/2003 Bastom University

LIB.

3

SECOND SEMESTER

(MARCH/APRIL 2004)

PH 304 CONDENSED STATE PHYSICS

You may assume that atomic mass of Almojokgo is 26.78e and the Arogadro's number is 6.023 × 10²² mole⁻¹

Time: 01 hour.

Answer <u>ALL</u> Questions

 (a) Define the terms lattice, basis, conventional unit cell of a crystal structure. What do you understand by Miller indices (hkl) of a crystal plane? Show that the spacing between consecutive parallel planes of Miller indices (hkl) in a cubic crystal of lattice constant a is given by

$$d_{hkl} = \frac{a}{\sqrt{h^2 + k^2 + l^2}}$$

- (b) What do you understand by *packing fraction* of a crystal structure? Crystal structure of a metal is *fcc*. The spacing d_{100} between adjacent (100) planes of the crystal is $2A^0$. Calculate
 - (i) radius of the atoms in the crystal
 - (ii) packing fraction of the crystal structure.

What are the assumptions you have made in these calculations?

- 2. (a) If the Aluminium lattice is fcc and the nearest neighbor distance between Aluminium atoms is $2.86A^0$. find
 - (i) The lattice constant
 - (ii) The spacing between (111) planes.
 - (iii) The number of atoms per unit volume
 - (iv) The density of Aluminium

You may assume that atomic mass of Aluminium is 26.78g and the Avogadro's number is $6.023 \times 10^{23} mole^{-1}$.

(b) Briefly describe the Bragg's diffraction in crystals and show that the Bragg condition for crystal diffraction on (hkl) planes is

$$2d_{hkl}sin\theta_{hkl} == n\lambda$$

The symbols have their usual meanings.

 $K_{\alpha} - X$ Rays from a target are incident on a cubic crystal of lattice constant 4.06 A^0 and the diffracted beam from (111) planes is observed at a Bragg angle 8.7⁰. Assuming order of diffraction n = 1 determine the wave length of $K_{\alpha} - X$ Ray.